



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**THE OCCURRENCE OF *BUFO COLUMBIENSIS* EAST OF THE ROCKY MOUNTAINS.**

BY ROBERT T. YOUNG.

High mountain chains, especially those whose crests extend above timber line, are more or less efficient barriers to the spread of animals. This is more particularly so in the case of species which do not migrate easily from place to place.

The capture of *Bufo columbiensis* on the eastern slope of the Continental Divide in Colorado is, therefore, of some interest.

In August, 1902, I obtained a specimen of this toad on the eastern slopes of the Arapahoe Peaks, about 28 km. west of Boulder, Colo., at an altitude of about 3,100 m., and it has since been reported to me by Prof. Beardsley, of the Colorado State Normal School, from the eastern slope of the mountains west of Greeley.

Cope<sup>1</sup> gives this species as a characteristic of the Pacific district of the Western sub-region of the Medicolumbian region, included in the Transition of Merriam.<sup>2</sup> My records, however, show it to occur in the Hudsonian zone, far above the upper limit of the Transition, where it has doubtless migrated from its original habitat.

There are several passes in northern Colorado over the Continental Divide which cross the mountains below timber line, affording possible paths of migration for toads from the western to the eastern slope.

Further observations on the distribution of this species on both slopes of the mountains, taken from year to year, would doubtless be of interest as affording some data for a determination of the conditions and rate of migration of toads.

---

<sup>1</sup> The Geographical Distribution of Batrachia and Reptilia in North America, *Am. Nat.*, XXX, 886-902 and 1003-1026.

<sup>2</sup> Laws of Temperature Control of the Geographic Distribution of Terrestrial Animals and Plants, *Nat. Geog. Mag.*, VI, 229-238.